



COVID-19 vaccination and blood clotting

Information about your vaccination

The UK vaccination programme has been very successful with more than 30 million people vaccinated and more than 6,000 lives already saved.

What is the concern?

Recently there have been reports of a very rare condition involving blood clots and unusual bleeding after vaccination. This is being carefully reviewed but the risk factors for this condition are not yet clear.

Although this condition remains extremely rare there appears to be a higher risk in people shortly after the first dose of the AstraZeneca (AZ) vaccine. Around 4 people

develop this condition for every million doses of AZ vaccine doses given.

This is seen slightly more often in younger people and tends to occur between 4 days and 2 weeks following vaccination.

This condition can also occur naturally, and clotting problems are a common complication of COVID-19 infection. An increased risk has not yet been seen after other COVID-19 vaccines but is being carefully monitored.

What are the benefits and risks of the vaccination?

Age	Risk from COVID-19	Benefit of vaccination	Risk of vaccination
Over 50 years of age or with underlying medical conditions	<p>Low</p> <ul style="list-style-type: none"> • catching infection • passing on infection <p>Very high</p> <ul style="list-style-type: none"> • hospitalisation • intensive care admission • death <p>Moderate</p> <ul style="list-style-type: none"> • Long COVID 	<p>One dose more than 80% reduction in:</p> <ul style="list-style-type: none"> • deaths • hospitalisation • intensive care <p>Two doses more than 95% reduction in:</p> <ul style="list-style-type: none"> • deaths 	<p>Uncommon</p> <ul style="list-style-type: none"> • sore arm • feeling tired • headache • general aches • flu like symptoms <p>Extremely rare</p> <ul style="list-style-type: none"> • clotting problems
30 to 49 years of age	<p>High</p> <ul style="list-style-type: none"> • catching mild infection • passing on infection <p>Low</p> <ul style="list-style-type: none"> • hospitalisation • intensive care admission • death <p>Moderate</p> <ul style="list-style-type: none"> • Long COVID 	<p>One dose between 60% and 70% reduction in:</p> <ul style="list-style-type: none"> • catching infection • passing on infection <p>Two doses more than 85% reduction in:</p> <ul style="list-style-type: none"> • catching and passing on infection 	<p>Common</p> <ul style="list-style-type: none"> • sore arm • feeling tired • headache • general aches • flu like symptoms <p>Extremely rare</p> <ul style="list-style-type: none"> • clotting problems
18 to 29 years of age	<p>Very high</p> <ul style="list-style-type: none"> • catching mild infection • passing on infection <p>Very low</p> <ul style="list-style-type: none"> • hospitalisation • intensive care admission • death <p>Moderate</p> <ul style="list-style-type: none"> • Long COVID 	<p>One dose between 60% and 70% reduction in:</p> <ul style="list-style-type: none"> • catching infection • passing on infection <p>Two doses more than 85% reduction in:</p> <ul style="list-style-type: none"> • catching and passing on infection 	<p>Very common</p> <ul style="list-style-type: none"> • sore arm • feeling tired • headache • general aches • flu like symptoms <p>Extremely rare</p> <ul style="list-style-type: none"> • clotting problems

What should I do next?

Over 50 years of age or with underlying medical conditions

All older adults (including health and social care workers over 50 years of age), care home residents, health and social care workers* and adults with certain medical conditions were prioritised in the first phase of the programme because they were at high risk of the complications of COVID-19.

The Medicines and Healthcare products Regulatory Agency (MHRA) and the Joint Committee on Vaccination and Immunisation (JCVI) advises that you should still receive any of the available COVID-19 vaccines. The benefits of vaccination in protecting you against the serious consequences of COVID-19 outweigh any risk of this rare condition. You should also complete your course with the same vaccine you had for the first dose.

If your first dose was with AZ vaccine without suffering any serious side effects you should have the second dose on time as you may still be at high risk of the complications of COVID-19. Having the second dose will give you higher and longer lasting protection.

If you are a healthy person over 30 to 50 years of age

The MHRA and the JCVI advises that all adults in this age group (including health and social care workers) should still receive any of the available COVID-19 vaccines. The benefits of vaccination in protecting you against the serious consequences of COVID-19 outweigh any risk of this rare condition. You should also complete your course with the same vaccine you had for the first dose.

If you are a healthy younger person aged 18 to 29

The MHRA and the JCVI continue to monitor the benefits and safety of the AZ vaccine in younger people. You should carefully consider the risk to both you and your family and friends of COVID-19 before making a decision.

Currently JCVI has advised that it is preferable for people under 30 to have a vaccine other than AZ. If you choose to have another COVID-19 vaccine you may have to wait to be protected. You may wish to go ahead with the AZ vaccination after you have considered all the risks and benefits for you.

*This includes unpaid carers and family members of those who are immunosuppressed

What should I look out for after vaccination?

Although serious side effects are very rare, if you experience any of the following from around 4 days to 4 weeks after vaccination you should seek medical advice urgently.

- a new, severe headache which is not helped by usual painkillers or is getting worse
- a headache which seems worse when lying down or bending over or
- an unusual headache that may be accompanied by:
 - blurred vision, nausea and vomiting
 - difficulty with your speech
 - weakness, drowsiness or seizures
- new, unexplained pinprick bruising or bleeding
- shortness of breath, chest pain, leg swelling or persistent abdominal pain

If you have any concerns after your vaccination about side effects, please speak to your GP.

For more information, visit [gov.je/vaccine](https://www.gov.je/vaccine)

What about the second dose?

The COVID-19 vaccine that you have had has been shown to reduce the chance of you suffering from COVID-19 disease. Each vaccine has been tested in more than 20,000 people in several different countries and shown to be safe.

It takes a few weeks for your body to build up protection from the vaccine. Like all medicines, no vaccine is completely effective, so you should continue to take recommended precautions to avoid infection. Some people may still get COVID-19 despite having a vaccination, but this should be less severe.